

Object oriented design and implementation of an intra-nuclear transport model

*L. Bellagamba¹, A. Brunengo², E. Di Salvo^{2,3}, M.G. Pia^{2,4}
for the Geant4 Collaboration*

¹ INFN, Sezione di Bologna, Italy

² INFN, Sezione di Genova, Italy

³ University of Genova, Italy

⁴ CERN, Switzerland

Abstract

The Hadron Kinetic Model is part of the Geant4 Toolkit; it covers the simulation of hadronic interactions in the intermediate energy range. Its role in the context of Geant4 hadronic models is outlined with emphasis on the physics content, and its Object Oriented design is illustrated.

Keywords: simulation, hadronic, Geant4

1 Introduction

The Hadron Kinetic Model is part of the set of theory driven models for hadronic interactions [1] in the Geant4 Simulation Toolkit [2].

The Hadron Kinetic Model covers the intermediate energy régime of hadronic interactions; it can satisfy two use cases for intranuclear transport, both as a hadronic interaction model to be used by the physics processes, and as a back-end to higher energy models.

Object Oriented technologies play a fundamental role to address the complex physics domain described by this model. The Object Oriented design also provides the basis for extensive code reuse by other models, such as cascade models or quantum molecular dynamics models; it also provides the means for easy extension or evolution of the system.

2 The physics of the Hadron Kinetic Model

Geant4 Hadron Kinetic Model implements the physics describing meson-meson, meson-baryon, baryon-baryon interactions, involving resonance excitation and deexcitation and particle absorption, in the energy range O(100 MeV) to O(5 GeV); it includes a large set of exclusive channels.

Due to the complexity of the physics handled by the Hadron Kinetic Model, only a few main features are outlined in this paper; a more detailed description is available in reference [6].

2.1 The Kinetic Algorithm

The Kinetic Algorithm can be considered a step by step updating of a particle vector; its implementation is performed according to the following scheme.

1. Create a vector of particles: assign initial particle types, their coordinates and momenta etc.; assign initial value for the time evolution parameter.
2. For a given step of the time evolution parameter find pairs of particles, according to a collision criterion, which are assumed to collide and particles which, according to their lifetimes, are assumed to decay.
3. Perform particle collisions and particle decays, determining the generation of outgoing particles; during this step particle coordinates and momenta are updated (particle propagation).

4. Starting from 2, perform the next step.

2.2 The scattering: cross sections and final states

The Hadron Kinetic Model handles two-body hadron elastic and inelastic scattering, including resonance excitation and deexcitation. The processes implemented cover baryon-baryon interactions (including baryon resonances), baryon-antibaryon annihilation, meson-baryon and meson-meson interactions.

The cross sections are functions of the incoming and outgoing particle types, their internal quantum numbers, like isospin, and energies. The overall physics approach of the Hadron Kinetic Model in this domain closely follows the one adopted by the UrQMD model [4]. The cross sections are calculated through various methods: from tabulations of experimental data, from parameterisations according to an algebraic function or from other cross sections via general principles, such as detailed balance [3], [4] or the additive quark model [5].

The angular distributions for all the relevant two-body processes are assumed to be similar and are determined according to the procedure described in reference [4].

Resonance decay channels are sampled according to the partial decay widths, which depend on the mass of the resonance.

Hadron-nucleus or nucleus-nucleus processes involve hadron-hadron collisions in high density nuclear matter. Under these extreme conditions the Pauli-blocking, which modifies the process vacuum behaviour, reducing a free particle cross section or decay width to an effective one due to Fermi statistics, must be taken into account. Each possible collision or decay is validated by checking the phase space fermion density, since a process can occur only if there is enough room (according to the Pauli principle) to accommodate the final state fermions. This effect is implemented treating fermions as gaussian shaped density distributions in the phase space. For each new fermion, produced by a possible collision or decay, a phase space overlapping density function is calculated and used to simulate Pauli-blocking. A detailed description of the used procedure is available in reference [6].

2.3 The field propagation

At every step, before performing an interaction - i.e. particles collision or particle decay - coordinates and momenta of all particles travelling through the nucleus are updated from the values at the time of previous interaction to the current time. The transportation of these particles is done describing the particle-nucleus interaction using phenomenological mean-field potentials, the so called “optical potentials” [6].

The particle propagation can be operated either with the cascade approach, i.e. along a straight line trajectory, using potentials only to evaluate the final particle energy at the end of the step, or with a numerical integration of the equations of motion, using the classical Runge-Kutta method.

3 The object oriented analysis and design

The Hadron Kinetic Model is integrated in the framework of Geant4 Theory Driven Hadronic Models [7].

The domain analysis has led to the identification of the main areas of responsibilities in the Model: the overall management, the handling of the scattering process and the propagation through the nuclear field. Additional utilities, such as the the description of the nucleus and of the particles involved in the interactions, are common to the whole set of Geant4 hadronic models.

Figure 1: An overview of the Hadron Kinetic Model design: top level class diagram.

The `G4HadronKineticModel` inherits from the `G4VIntraNuclearTransportModel` class, that fulfills the role of providing to the intra-nuclear transport models an interface to the processes as well as the possibility to be used as a back-end to high energy models. The `G4HadronKineticModel` class has the responsibility of managing the main functionality of the model, as described in section 2.1. It delegates the responsibility of the scattering and of the field transport to the `G4VScatterer` and `G4VFieldPropagation` classes respectively.

The concrete class `G4Scatterer`, inheriting from `G4VScatterer`, handles the scattering process. It manages the complex physics content of the scattering process in the energy domain of the intra-nuclear transport, including the resonance region, through a design based on a recurrent Composite Pattern [8]. The scattering channel classes inherit from the `G4VCollision` abstract base class; some of them are composed of sub-channels, modeled through the Composite Pattern (for instance, the “ExcitedNucleon-Nucleon” scattering channel consists of sub-channels such as `N1440-p`, `N1520-n` etc.). The Composite Pattern also handles the specialization of collisions into elastic (leaf) and inelastic (composite) ones. The Composite Pattern is also adopted to handle the cross section sector, mapping the availability of experimental data or theoretical cross section formulae for the collisions: for instance, the total cross section of a channel is represented by a leaf cross section whenever specific data or model are available and by a composite cross section (composed of single branch cross sections) when no total measurements or models are available; both leaf and composite cross sections are handled transparently by the Collisions through the same abstract interface to `G4VCrossSectionSource`.

The Strategy Pattern [8] is used throughout the design to handle interchangeable algorithms, whenever alternative models or parameterisations for the same physics item are available.

This flexible design of the Scatterer allows for an easy extensibility of the set of scattering processes handled by the Hadron Kinetic Model, whenever new theoretical descriptions or experimental data become available to improve the overall precision of the physics of the model itself; at the same time, it makes alternative physics implementations for single scattering channels easily interchangeable, thus allowing to investigate and optimise the physics capabilities of the model. The clear separation between the cross section sector and the generation of final states further enhances the flexibility of the Scatterer to accommodate alternative modelling. The recurrent Composite Pattern contributes to improve the overall performance, by enabling to establish a logical hierarchy in the messaging of the physics components. The general design of the Scatterer makes it suitable to be used also by other hadronic simulation models (such as, for instance, the QMD Model or cascade models), as well as in other contexts of Geant4 physics simulation.

The field propagation is handled by the `G4VFieldPropagation` class, that is virtual to allow different strategies to be used. The concrete `G4RKFieldIntegrator` class derives from it, and performs the propagation of particles via a Runge-Kutta numerical integration. It uses classes for optical potentials - `G4VKMOpticalField` - and for their derivatives - `G4VKMOpticalEqRhs` - that are made virtual to implement again a Strategy Pattern. `G4VKMOpticalEqRhs` inherits from the `G4MagEqRhs` class; this allows to reuse in the Kinetic Model the code common to the Geometry-Transportation domain for the integration of the equations of motion.

The Hadron Kinetic Model makes use of other design components common to the Geant4 Hadronic Models, such as the classes responsible for the description of the nucleus and of the particles involved in the interactions.

4 Conclusions

The Geant4 Hadron Kinetic Model represents a novel approach to intra-nuclear transport. Object Oriented methodologies have been a key factor to handle the complexity of the underlying physics and to ensure the future extensibility and maintenance of the model.

At the time of the CHEP2000 Conference a first implementation of the Hadron Kinetic Model is available in Geant4 and is under thorough software and physics testing. It is meant to become part of the Geant4 public release, following the completion of the testing phase.

References

- 1 The Geant4 Collaboration, Geant4 Physics Reference Manual, <http://wwwinfo.cern.ch/asd/geant4/G4UsersDocuments/UsersGuides/PhysicsReferenceManual/html/PhysicsReferenceManual.html>, 1999.
- 2 The RD44 Collaboration, CERN/LHCC 98-44, 1998.
The Geant4 Collaboration, <http://wwwinfo.cern.ch/asd/geant4/geant4.html>.
- 3 P. Danielewicz and G.F. Bertsch, Nucl. Phys. A533 (1991) 712.
- 4 A.S. Bass et al., Prog. Part. Nucl. Phys. 41 (1998) 225.
- 5 H.J. Lipkin and F. Scheck, Phys. Rev. Lett. 16 (1966) 71.
J.J.J. Kokkede, The Quark Model, W.A. Benjamin Inc., New York, 1969.
- 6 L. Bellagamba et al., Simulation of interactions of hadrons with the hadronic matter: the Hadron Kinetic Model, INFN/AE preprint, to be submitted.
- 7 J.P. Wellisch for the Geant4 Collaboration, Proceedings of CHEP98, Chicago, 1998.
- 8 E. Gamma et al., Design Patterns, Addison-Wesley, 1994.